Settings	Clinical Areas	Recommended use of PPE				
		Yellow	Frequency of PPE use	Orange	Frequency of PPE use	
Emergency Department	Screening/ Nurse triage assessment	N95	Extended use up to 6 hrs [^]	N95 mask	Extended use up to 6 hrs [^]	
	Fever area	Full PPE: N95 mask, gown, gloves, eye protection (if available)	Extended use up to 6 hrs [^] for masks & eye protection (if used). Single use for gown and gloves, i.e. change between patients	Full PPE: N95 mask, gown, gloves, eye protection (if available)	Extended use up to 6 hrs [^] for masks and eye protection (if used). Single use for gown and gloves	
	All other areas	Surgical mask	Extended use up to 6 hrs [^]	Surgical mask	Extended use up to 6 hrs [^]	
Isolation ward	Rooms with suspected/ confirmed cases	Full PPE: N95 mask, gown, gloves, eye protection	Extended use up to 6 hrs [^] for eye protection. Single use for N95, gown, gloves.	Full PPE: N95 mask, gown, gloves, eye protection	Extended use up to 6 hrs [^] for mask and eye protection. Single use for gown and gloves.	
	Non-clinical staff (not entering isolation rooms)	n/a	n/a	Surgical mask	Extended use up to 6 hrs [^]	
	All other areas	No mask required; mask if droplet precaution needed	-	Surgical mask	Extended use up to 6 hrs [^]	
ICU	Rooms with suspected/ confirmed cases	Full PPE: N95 mask, gown, gloves, eye protection	Extended use up to 6 hrs for eye protection. Single use for N95 mask, gown, gloves.	Full PPE: N95 mask, gown, gloves, eye protection	Extended use up to 6 hrs [^] for mask & eye protection. Single use for gown, gloves.	
	Rooms without suspected/ confirmed cases	As per peacetime requirement	-	Surgical mask	Extended use up to 6 hrs [^]	
	All other areas	No mask required; surgical mask if droplet precaution needed	-	Surgical mask	Extended use up to 6 hrs [^]	

Settings	Clinical Areas	Recommended use of PPE				
		Yellow	Frequency of PPE use	Orange	Frequency of PPE use	
General ward, Community Hospital	Rooms with clinically vulnerable patients (e.g. transplant/ renal patients)	Surgical mask	Single use	Surgical mask	Single use	
	All other areas	No mask required; surgical mask if droplet precaution needed	-	Surgical mask	Extended use up to 6 hrs [^]	
SOCs [#] , other	Triage staff	Surgical mask	Extended use up to 6 hrs [^]	Surgical mask	Extended use up to 6 hrs [^]	
ambulatory services e.g. Pharmacy, Radiology, Physio, Dental Clinics, other Specialist Clinics	All other areas	No mask required; surgical mask if droplet precaution needed	-	Surgical mask	Extended use up to 6 hrs [^]	
All other non- clinical areas [®]	Administrative, offices, store rooms, etc. where there is no direct patient contact	No PPE required	-	No PPE required	-	
Polyclinics	Triage staff	Surgical mask	Extended use up to 6 hrs	N95 mask	Extended use up to 6 hrs [^]	
	All other areas	No mask required; surgical mask if droplet precaution needed	-	Surgical mask	Extended use up to 6 hrs [^]	
	Routine cases (e.g. general consultation)	No mask required; surgical mask if droplet precaution needed	-	Surgical mask	Extended use up to 6 hrs [^]	
	Suspected cases (Segregation/ isolation room)	N95 mask, gown, gloves	Single use	N95 mask, gown, gloves	Extended use up to 6 hrs [^] for mask. Single use for gown, gloves.	

Settings	Clinical Areas	Recommended use of PPE				
		Yellow	Frequency of PPE use	Orange	Frequency of PPE use	
GP Clinics, Community paediatricians	Triage staff	Surgical mask	Extended use up to 6 hrs	N95 mask	Extended use up to 6 hrs ²	
	Routine cases (e.g. general consultation)	No mask required; surgical mask if droplet precaution needed	-	Surgical mask	Extended use up to 6 hrs ²	
	Suspected cases	N95 mask, gown, gloves	Single use	N95 mask, gown, gloves	Extended use up to 6 hrs ² for mask. Single use for gown, gloves	
General ward, Community Hospital	Rooms with clinically vulnerable patients (e.g. transplant/ renal patients)	Surgical mask	Single use	Surgical mask	Single use	
	All other areas	No mask required; surgical mask if droplet precaution needed	-	Surgical mask	Extended use up to 6 hrs [^]	
Community- based care e.g. Senior Activity Centres	Routine care	As per peacetime requirement	-	Surgical mask	Extended use up to 6 hrs [^]	
Home-based care	Routine care	As per peacetime requirement	-	Surgical mask	Extended use up to 6 hrs [^]	

Footnotes:

[^]Extended use should be for the period when staff are in clinical areas (exceptions include evident contamination of PPE e.g. blood spillage, soiled, or soggy, in which case staff should change PPEs). Staff must strictly adhere to hand hygiene practices in addition to PPE use to prevent cross contamination.

* SOC/ Dental/ other private specialist clinic attendances should be provided screening questions for travel history and respiratory symptoms, and these cases should be referred to ED/ GP clinics for further triage if the patient fulfils criteria for suspect case or when assessed to be of high risk by a clinician.

[®]Laboratories are to continue with their own current PPE measures