Recommendations on the Use of the Sinovac-CoronaVac Vaccine

S/N	Recommended Use of the Sinovac-CoronaVac Vaccine
1	All persons are strongly recommended to get PSAR-authorized mRNA vaccines for their primary vaccine series, unless medically ineligible.
	Those who are recommended to get a booster, after completing their primary vaccine series, should get mRNA vaccines as their booster (unless unable for medical reasons such as allergy or myocarditis from a prior dose).
2	For persons who are unable to take a second dose of mRNA vaccines (M) for medical reasons, they should complete their vaccination with the Sinovac-CoronaVac vaccine (S):
	M-S-S If medically unable to take second dose of mRNA vaccine, recommend 2 doses of Sinovac-CoronaVac. Dose 2 is 28 days after Dose 1.
3	For persons who already completed their primary series with mRNA vaccines (M) and are recommended to get a booster, but are unable to take another dose of mRNA vaccines (M) for medical reasons, they should boost using one dose of Sinovac-CoronaVac vaccine (S):
	M-M + S If medically unable to take additional doses of mRNA vaccine, recommend one dose of Sinovac-CoronaVac if booster is needed. Booster is 6 months after mRNA Dose 2; denoted with suffix (+).
4	For persons who have already received two doses of Sinovac-CoronaVac vaccine (S), and can take mRNA vaccines (M), they are recommended to get an mRNA vaccine (M) to complete their primary vaccination series:
	S-S-M One dose of mRNA vaccine is recommended, given at 90 days after Sinovac Dose 2 to complete the primary series.
5	<u>F</u> or persons who have already received two doses of Sinovac-CoronaVac vaccine (S), and can take mRNA vaccines (M) but decline for personal reasons, they should complete the primary series with a third dose, as follows:
	S-S-S Dose 3 at 90 days after Dose 2 to complete the primary series.